


Baby it's Hot in Here: Impacts of Late Gestation Heat Stress on Dam and Daughter

G. E. Dahl




Department of Animal Sciences
 Institute of Food and Agricultural Sciences
gdahl@ufl.edu
 Balchem Real Science Series
 4 May 2021









Gainesville, Florida, USA

- Sand bedded free stalls
- Fans over stalls
- Soakers over feedline
- Fans on at 70° F (21.1°C)
- Soakers on 1.5 min every 5 min at 72° F










Late Gestation Cooling


- Do I have to cool cows the entire dry period?
- Do heifers need to be cooled pre-partum?

Heat Stress Increases Rectal Temperature and Respiration Rate






Fabris et al., *J. Dairy Sci.* 102:5647-5656.


Heat Stress at Any Time Decreases Gestation Length and Dry Period Length




Fabris et al., *J. Dairy Sci.* 102:5647-5656.




Heat Stress Reduces DMI Prepartum But Not Postpartum




Tao et al., *J. Dairy Sci.* 94:5976–5986

Cooling Dry Cows Increases BW Prepartum, Decreases Postpartum

Thompson et al., *J. Dairy Sci.* 97:7426–7436

Effect of Cooling Dry Cows on Metabolic Profile

Tao et al., *J. Dairy Sci.* 95:5035–5046



Dry in COOL Months Improves Performance

Table 1. Milk production and occurrence of mastitis, digestive and respiratory problems, retained fetal membranes, and metritis in cows dried during HOT months (Jun, Jul, Aug) or COOL months (Dec, Jan, Feb) in the first 80 DIM of the subsequent lactation

Item	Dry during HOT months (Jun, Jul, Aug), n = 1,569				Dry during COOL months (Dec, Jan, Feb), n = 1,044				P-value
	Value	Disease ¹	n	%	Value	Disease ¹	n	%	
Milk production (kg)	10,351 ± 59.8				10,902 ± 73.3				0.01
Mastitis	0	1,288	82.0		0	650	91.0	0.01	
	1	283	18.0		1	94	9.0		
Digestive	0	1,516	98.6		0	973	93.2	0.01	
	1	53	3.4		1	71	6.8		
Respiratory	0	1,346	85.8		0	942	90.2	0.01	
	1	223	14.2		1	102	9.8		
Retained fetal membranes	0	1,586	99.6		0	1,100	97.0	0.06	
	1	69	4.4		1	31	3.0		
Metritis	0	1,500	95.6		0	1,007	98.4	0.35	
	1	67	4.2		1	38	3.5		

¹Disease: 0 = cows without the disease; 1 = cows with the disease.


Thompson & Dahl, *Prof. Anim. Sci.* 28:628-631




Dry in COOL Months Improves Reproductive Performance

Table 3. Milk production and reproductive performance of cows dried during HOT months (Jun, Jul, Aug) or COOL months (Dec, Jan, Feb) in the first 150 DIM of the subsequent lactation on a commercial farm in Florida


Item	Dry during HOT months (Jun, Jul, Aug)	Dry during COOL months (Dec, Jan, Feb)	P-value
Milk production (kg)	10,547 ± 67.0	11,005 ± 83.38	0.01
Number of breedings (n)	1,048	676	0.03
Mean (no.)	1.59 ± 0.02	1.51 ± 0.03	
DIM to breeding (n)	1,047	676	0.01
Mean (d)	91.0 ± 0.74	91.9 ± 0.92	
DIM to pregnancy (n)	1,051	679	0.01
Mean (d)	131.1 ± 0.85	125.9 ± 1.06	

Thompson & Dahl, *Prof. Anim. Sci.* 28:628-631






Why Does Cooling Affect AEA? Calf or Colostrum Effect?




Monteiro et al., J. Dairy Sci. 97:6426-6439

Experiment 1
- In utero
heat stress
for ~6 weeks
reduces body
weight and
height to
weaning



Monteiro et al., J. Dairy Sci. 97:6426-6439

Cooling Increased Apparent Efficiency of IgG Absorption (AEA*)



* AEA = [Serum [IgG] (g/L) * birth weight (kg) * 0.091 / IgG fed (g)] x 100

Monteiro et al., J. Dairy Sci. 97:6426-6439

Experiment 2 – No Effect of Colostrum from Cooled or Heat Stressed Cows on Calf Performance

Growth performance of calves born to cows under thermoneutral conditions during the dry period and fed frozen colostrum from cows exposed to either heat stress or cooling during the dry period

Parameter	Heat Stress LSM ± SE	Cooling LSM ± SE	P-value
Birth Weight (kg)	38.8 ± 1.4	39.2 ± 1.5	0.8
Weaning Weight (kg) ¹	68.4 ± 2.5	64.8 ± 2.6	0.4
Preweaning BW Gain (kg) ²	29.6 ± 2.3	25.6 ± 2.4	0.3
Avg. Daily Gain (kg/d)	0.49 ± 0.7	0.43 ± 0.8	0.2
Weaning Withers Height (cm) ¹	84.3 ± 0.8	83.0 ± 0.9	0.4
Preweaning Height Increase (cm) ²	7.8 ± 1.1	6.2 ± 1.0	0.3

¹Weaning weight and weaning height were measured at d 60 of age.

²Preweaning BW gain and height increase was calculated by individually subtracting data at d 60 of age by data at birth.

Monteiro et al., *J. Dairy Sci.* 97:6426–6439

J. Dairy Sci. 92:5988–5999

doi:10.3168/jds.2009-2243

© American Dairy Science Association, 2009

Heat stress abatement during the dry period:

Does cooling improve transition into lactation?

B. C. do Amaral,¹ E. E. Connor,² S. Tao,³ J. Hayen,⁴ J. W. Bullock,⁵ and G. E. Dahl⁶

¹Department of Animal Sciences, University of Florida, Gainesville, 32611

²USDA-ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705

³USDA-ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705

⁴USDA-ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705

⁵USDA-ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705

⁶University of Florida, Institute of Food and Agricultural Sciences, Gainesville, FL 32611

Heat stress abatement during the dry period influences

metabolic gene expression and improves immune

status in the transition period of dairy cows

B. C. do Amaral,¹ E. E. Connor,² S. Tao,³ J. Hayen,⁴ J. W. Bullock,⁵ and G. E. Dahl⁶

¹Department of Animal Sciences, University of Florida, Gainesville, 32611

²USDA-ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705

³USDA-ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705

⁴USDA-ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705

⁵USDA-ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705

⁶University of Florida, Institute of Food and Agricultural Sciences, Gainesville, FL 32611

Effect of heat stress during the dry period on mammary gland development

S. Tao,¹ J. W. Bullock,² B. C. do Amaral,¹ I. M. Thompson,¹ M. J. Hayen,³ S. E. Johnson,⁴ and G. E. Dahl⁵

¹Department of Animal Sciences, University of Florida, Gainesville, 32611

²USDA-ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705

³USDA-ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705

⁴USDA-ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705

⁵University of Florida, Institute of Food and Agricultural Sciences, Gainesville, FL 32611

Effect of cooling heat-stressed dairy cows during

the dry period on insulin response

S. Tao,¹ I. M. Thompson,¹ A. P. A. Monteiro,² M. J. Hayen,³ L. J. Young,⁴ and G. E. Dahl⁵

¹Department of Animal Sciences, University of Florida, Gainesville, 32611

²USDA-ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705

³USDA-ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705

⁴USDA-ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705

⁵University of Florida, Institute of Food and Agricultural Sciences, Gainesville, FL 32611

Effect of cooling during the dry period on immune response after

Streptococcus uberis intramammary infection challenge of dairy cows

I. M. T. Thompson,¹ S. Tao,² A. P. A. Monteiro,² K. C. Jeong,³ and G. E. Dahl⁴

¹Department of Animal Sciences, University of Florida, Gainesville, 32611

Retrospective analysis of records of calves from 5 studies between 2007 and 2011

Monteiro et al., *J. Dairy Sci.* 99:8443–8450

J. Dairy Sci. 92:5988–5999

doi:10.3168/jds.2009-2243

© American Dairy Science Association, 2009

Heat stress abatement during the dry period:

Does cooling improve transition into lactation?

B. C. do Amaral,¹ E. E. Connor,² S. Tao,³ J. Hayen,⁴ J. W. Bullock,⁵ and G. E. Dahl⁶

¹Department of Animal Sciences, University of Florida, Gainesville, 32611

²USDA-ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705

³USDA-ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705

⁴USDA-ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705

⁵USDA-ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705

⁶University of Florida, Institute of Food and Agricultural Sciences, Gainesville, FL 32611

Heat Stress Experiments 2007 - 2011

	Bulls	Heifers	Total
Cooling	31	41	72
Heat Stress	30	44	74
Total	61	85	147

J. Dairy Sci. 97:7426–7435

http://dx.doi.org/10.3168/jds.2013-7621

© American Dairy Science Association[®], 2014

Effect of cooling during the dry period on immune response after

Streptococcus uberis intramammary infection challenge of dairy cows




I. M. T. Thompson,¹ S. Tao,² A. P. A. Monteiro,² K. C. Jeong,³ and G. E. Dahl⁴

¹Department of Animal Sciences, University of Florida, Gainesville, 32611

²USDA-ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705

³USDA-ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705

⁴University of Florida, Institute of Food and Agricultural Sciences, Gainesville, FL 32611



In Utero Heat Stress Decreases Reproductive Performance


Table 2. Effect of maternal heat stress (HT) or cooling (CL) during late gestation on reproductive performance before first lactation of heifers born to HT or CL dams

Parameter	CL	HT	SEM	P
N	36	32	---	---
Age at first AI, mo	13.6	13.8	0.2	0.32
Services per pregnancy d ¹ 30	2.0	2.5	0.2	0.05
Age at pregnancy d ¹ 30, mo	16.1	16.9	0.3	0.07
Services per pregnancy d ¹ 50	2.3	2.6	0.2	0.32
Age at calving, mo	24.8	25.0	0.4	0.72

¹Days after insemination.


Monteiro et al., *J. Dairy Sci.* 99:8443-8450.

In Utero Heat Stress Reduces Milk Production



Monteiro et al., *J. Dairy Sci.* 99:8443-8450.

In Utero Heat Stress Does Not Affect Mature Bodyweight



Monteiro et al., *J. Dairy Sci.* 99:8443-8450.

